arXiv:1510.07778v2 [math.AT] 11 Mar 2016

MOMENT-ANGLE MANIFOLDS, 2-TRUNCATED CUBES AND MASSEY OPERATIONS

IVAN LIMONCHENKO

Abstract. We construct a family of manifolds having a nontrivial Massey n-product in their cohomology for any given n. These manifolds turn out to be smooth closed 2-connected manifolds with a compact torus $\mathbb{T}^m$-action called moment-angle manifolds $\mathcal{Z}_P$, whose orbit spaces are simple n-dimensional polytopes P obtained from a n-cube by a sequence of truncations of faces of codimension 2 only (*2-truncated cubes*). Moreover, the polytopes P are flag nestohedra but not graph-associahedra. For the latter class of flag polytopes Q we prove a formula for $\beta^{-i,2(i+1)}(Q)$ and relate it to the structure of the loop homology (Pontryagin algebra) $H_*(\Omega\mathcal{Z}_Q)$. We then study higher Massey products in $H^*(\mathcal{Z}_Q)$ for a graph-associahedron Q.

1. Introduction

The main aim of this work is to show that one of the key objects of study in toric topology – the moment-angle manifold $\mathcal{Z}_P$ of a simple convex n-dimensional polytope P – gives us an example of a smooth closed 2-connected manifold with a compact torus action such that its rational cohomology ring may contain a nontrivial higher Massey product of order n. These polytopes P are 2-truncated cubes and, moreover, flag nestohedra. This class of flag simple polytopes was studied in toric topology by V.M.Buchstaber and V.D.Volodin, who proved that flag nestohedra can be realized as 2-truncated cubes and that Gal conjecture on γ-vectors of simple polytopes holds for 2-truncated cubes and, thus, for all flag nestohedra [6]. In the polytope case we generalize the result of I.V.Baskakov [2] who constructed a family of simplicial complexes K whose moment-angle complexes $\mathcal{Z}_K$ have a non-trivial triple Massey product of 3-dimensional classes in $H^*(\mathcal{Z}_K)$. In the lowest dimension Baskakov construction gives a 2-sphere with 8 vertices K – the only K with a nontrivial triple Massey product in $H^*(\mathcal{Z}_K)$ among all the fourteen 2-spheres on 8 vertices. G.Denham and A.Suciu [8] generalized the result of Baskakov by proving a combinatorial criterion for K to give a $\mathcal{Z}_K$ with a non-trivial triple Massey product of 3-dimensional classes in $H^*(\mathcal{Z}_K)$.

Denote by K a simplicial complex of dimension $n-1$ on the vertex set $[m]=\{1,\dots,m\}$ and by $\Bbbk$ the base field or the ring of integers. Let $\Bbbk[v_1,\dots,v_m]$ be the graded polynomial algebra on m variables, $\deg(v_i)=2$.

2010 *Mathematics Subject Classification.* Primary 13F55, 55S30, Secondary 52B11.

Key words and phrases. moment-angle manifold, flag nestohedra, Stanley–Reisner ring, Massey products, graph-associahedron.

The work was supported by the RFBR grant no. 14-01-00537a.

The *Stanley–Reisner ring* (or the *face ring*) of K over $\Bbbk$ is the quotient ring

$$\Bbbk[K] = \Bbbk[v_1, \dots, v_m]/\mathcal{I}_K$$

where $\mathcal{I}_K$ is the ideal generated by square free monomials $v_{i_1} \cdots v_{i_k}$ such that $\{i_1, \dots, i_k\}$ is not a simplex in K. The monomial ideal $\mathcal{I}_K$ is called the *Stanley–Reisner ideal* of K. Then $\Bbbk[K]$ has a structure of a $\Bbbk$-algebra and a module over $\Bbbk[v_1, \dots, v_m]$ via the quotient projection.

In what follows we denote by P a simple n-dimensional convex polytope with m *facets* (i.e faces of codimension 1) $F_1, \dots, F_m$. Such a polytope P can be defined as a bounded intersection of m halfspaces:

$$P = \big\{\boldsymbol{x} \in \mathbb{R}^n \colon \langle \boldsymbol{a}_i, \boldsymbol{x}\rangle + b_i \geqslant 0 \quad \text{for } i = 1, \dots, m\big\}, \qquad (*)$$

where $\boldsymbol{a}_i \in \mathbb{R}^n$, $b_i \in \mathbb{R}$. We assume that the hyperplanes defined by the equations $\langle \boldsymbol{a}_i, \boldsymbol{x}\rangle + b_i = 0$ are in general position, that is, at most n of them meet at a single point. We also assume that there are no redundant inequalities in $(*)$, that is, no inequality can be removed from $(*)$ without changing P. Then the *facets* of P are given by

$$F_i = \big\{\boldsymbol{x} \in P \colon \langle \boldsymbol{a}_i, \boldsymbol{x}\rangle + b_i = 0\big\}, \quad \text{for } i = 1, \dots, m.$$

Let A_P be the $m \times n$ matrix of row vectors $\boldsymbol{a}_i$, and denote by $\boldsymbol{b}_P$ be the column vector of scalars $b_i \in \mathbb{R}$. Then we can rewrite $(*)$ as

$$P = \big\{\boldsymbol{x} \in \mathbb{R}^n \colon A_P\boldsymbol{x} + \boldsymbol{b}_P \geqslant \boldsymbol{0}\big\}.$$

Consider the affine map

$$i_P \colon \mathbb{R}^n \to \mathbb{R}^m, \quad i_P(\boldsymbol{x}) = A_P\boldsymbol{x} + \boldsymbol{b}_P.$$

which embeds P into

$$\mathbb{R}^m_{\geqslant} = \{\boldsymbol{y} \in \mathbb{R}^m \colon y_i \geqslant 0 \quad \text{for } i = 1, \dots, m\}.$$

Definition. We define the space $\mathcal{Z}_P$ as a pull-back in the following commutative diagram [5, Lemma 3.1.6, Construction 3.1.8]:

$$\begin{array}{ccc} \mathcal{Z}_P & \xrightarrow{\;i_Z\;} & \mathbb{C}^m \\ \big\downarrow & & \big\downarrow{\scriptstyle \mu} \\ P & \xrightarrow{\;i_P\;} & \mathbb{R}^m_{\geqslant} \end{array}$$

where $\mu(z_1, \dots, z_m) = (|z_1|^2, \dots, |z_m|^2)$. The latter map may be thought of as the quotient map for the coordinatewise action of the standard torus

$$\mathbb{T}^m = \{\boldsymbol{z} \in \mathbb{C}^m \colon |z_i| = 1 \quad \text{for } i = 1, \dots, m\}$$

on $\mathbb{C}^m$. Therefore, $\mathbb{T}^m$ acts on $\mathcal{Z}_P$ with quotient P, i_Z is a $\mathbb{T}^m$-equivariant embedding with a trivial normal bundle and $\mathcal{Z}_P$ is embedded into $\mathbb{C}^m$ as a nondegenerate intersection of Hermitian quadrics. One can easily see that $\mathcal{Z}_P$ has a structure of a smooth closed manifold of dimension $m + n$, called the *moment-angle manifold* corresponding to P.

Suppose $(\mathbf{X}, \mathbf{A}) = \{(X_i, A_i)\}_{i=1}^m$ is a set of topological pairs. The following notion appeared firstly in the work by V.M.Buchstaber and T.E.Panov [5] and then was studied intensively and generalized in the works of A.Bahri, M.Bendersky, F.Cohen, S.Gitler [1], J.Grbić and S.Theriault [11], K.Iriye and D.Kishimoto [13] and others.

Definition. A *polyhedral product* is a topological space:

$$(\mathbf{X},\mathbf{A})^K=\bigcup_{I\in K}(\mathbf{X},\mathbf{A})^I,$$

where $(\mathbf{X},\mathbf{A})^I=\prod_{i=1}^m Y_i$ for $Y_i=X_i$, if $i\in I$, and $Y_i=A_i$, if $i\notin I$. Particular cases of a polyhedral product $(\mathbf{X},\mathbf{A})^K$ include *moment-angle complexes* $\mathcal{Z}_K=(\mathbb{D}^2,\mathbb{S}^1)^K$ and *real moment-angle complexes* $\mathcal{R}_K=(\mathbb{D}^1,\mathbb{S}^0)^K$.

Denote by K_P the boundary ∂P^* of the dual simplicial polytope. It can be viewed as a $(n-1)$-dimensional simplicial complex on the set $[m]$, whose simplices are subsets $\{i_1,\dots,i_k\}$ such that $F_{i_1}\cap\ldots\cap F_{i_k}\neq\varnothing$ in P. By [4, Theorem 6.2.4], $\mathcal{Z}_P$ is $\mathbb{T}^m$-equivariantly homeomorphic to the moment-angle complex $\mathcal{Z}_{K_P}$.

The Tor-groups of K acquire a topological interpretation by means of the following result due to Buchstaber and Panov.

Theorem 1.1 ([4, Theorem 4.5.4] or [16, Theorem 4.7])**.** *The cohomology algebra of the moment-angle complex $\mathcal{Z}_K$ is given by the isomorphisms*

$$\begin{aligned}H^{*,*}(\mathcal{Z}_K;\Bbbk)&\cong\operatorname{Tor}^{*,*}_{\Bbbk[v_1,\dots,v_m]}(\Bbbk[K],\Bbbk)\\&\cong H\big[\Lambda[u_1,\dots,u_m]\otimes\Bbbk[K],d\big]\\&\cong\bigoplus_{I\subset[m]}\widetilde{H}^*(K_I;\Bbbk),\end{aligned}$$

where bigrading and differential in the cohomology of the differential bigraded algebra are defined by

$$\operatorname{bideg}u_i=(-1,2),\ \operatorname{bideg}v_i=(0,2);\quad du_i=v_i,\ dv_i=0.$$

In the third row, $\widetilde{H}^*(K_I)$ denotes the reduced simplicial cohomology of the induced subcomplex K_I of K (the restriction of K to $I\subset[m]$). The last isomorphism is the sum of isomorphisms

$$H^p(\mathcal{Z}_K)\cong\sum_{I\subset[m]}\widetilde{H}^{p-|I|-1}(K_I),$$

and the ring structure is given by the maps

$$\widetilde{H}^{p-|I|-1}(K_I)\otimes\widetilde{H}^{q-|J|-1}(K_J)\to\widetilde{H}^{p+q-|I|-|J|-1}(K_{I\cup J}),\qquad(*)$$

which are induced by the canonical simplicial maps $K_{I\cup J}\hookrightarrow K_I*K_J$ (join of simplicial complexes) for $I\cap J=\varnothing$ and zero otherwise.

Additively the following theorem due to M.Hochster holds.

Theorem 1.2 ([12])**.** *For any simplicial complex K on m vertices we have:*

$$\operatorname{Tor}^{-i,2j}_{\Bbbk[v_1,\dots,v_m]}(\Bbbk[K],\Bbbk)\cong\bigoplus_{J\subset[m],\,|J|=j}\widetilde{H}^{j-i-1}(K_J;\Bbbk).$$

The ranks of the bigraded components of the Tor-algebra

$$\beta^{-i,2j}(\Bbbk[K])=\operatorname{rk}_\Bbbk\operatorname{Tor}^{-i,2j}_{\Bbbk[v_1,\dots,v_m]}(\Bbbk[K],\Bbbk)$$

are called the *bigraded Betti numbers* of $\Bbbk[K]$ or K, when $\Bbbk$ is fixed. A particular case of the Hochster result is the following one. If $j = i+1$ then

$$\beta^{-i,2(i+1)}(P) = \sum_{J\subset[m],|J|=i+1} (\mathrm{cc}(P_J) - 1),$$

where $P_J = \cup_{j\in J} F_j$ and $\mathrm{cc}(P_J)$ equals the number of connected components of P_J.

Due to [4, Construction 3.2.8, Theorem 3.2.9] the Tor-algebra of K acqures a multigrading and the multigraded components can be calculated in terms of induced subcomplexes.

Theorem 1.3. *For any simplicial complex K on m vertices we have:*

$$\mathrm{Tor}^{-i,2J}_{\Bbbk[v_1,\ldots,v_m]}(\Bbbk[K],\Bbbk) \cong \widetilde{H}^{|J|-i-1}(K_J;\Bbbk),$$

where $J\subset[m]$ and $\mathrm{Tor}^{-i,2\mathbf{a}}_{\Bbbk[v_1,\ldots,v_m]}(\Bbbk[K],\Bbbk) = 0$, if $\mathbf{a}$ is not a $(0,1)$-vector.

Moreover, if we denote by $R(K) = \Lambda[u_1,\ldots,u_m]\otimes\Bbbk[K]/(v_i^2 = u_iv_i = 0, 1\leq i\leq m)$ a graded algebra with the differential d as in Theorem 1.1 then $R(K)$ also acquires multigrading and the following isomorphism holds:

$$\mathrm{Tor}^{-i,2\mathbf{a}}_{\Bbbk[v_1,\ldots,v_m]}(\Bbbk[K],\Bbbk) \cong H^{-i,2\mathbf{a}}[R(K),d]$$

for any simplicial complex K.

The author is grateful to Victor Buchstaber and Taras Panov for many helpful discussions and advice. I also thank Jim Stasheff for his comments and suggestions on improving the text.

2. Nestohedra and graph-associahedra

We begin with a definition of a family of simple polytopes called nestohedra and state the result of Buchstaber and Volodin on geometric realization of flag nestohedra.

Definition. Let $[n+1] = \{1,2,\ldots,n+1\}$, $n\geq 2$. *A* building set on [n+1] is a family of nonempty subsets $B = \{S\subseteq[n+1]\}$, such that: 1) $\{i\}\in B$ for all $1\leq i\leq n+1$, 2) if $S_1\cap S_2\neq\varnothing$, then $S_1\cup S_2\in B$. A building set is *connected* if $[n+1]\in B$.

Then a *nestohedron* is a simple convex n-dimensional polytope $P_B = \sum\limits_{S\in B}\Delta_S$, where in the Minkowski sum one has

$$\Delta_S = \mathrm{conv}\{e_j |\, j\in S\}\subset\mathbb{R}^{n+1}.$$

Note that facets of P_B are in 1-1 correspondence with proper elements S in B.

Example 2.1. If P is a combinatorial n-simplex then the subset of $2^{[n+1]}$ consisting of all the singletons $\{i\}, 1\leq i\leq n+1$ and the whole set $[n+1]$ gives a building set B, such that $P = P_B$ for any $n\geq 2$.
If P is a combinatorial n-cube then the following set B consisting of

$$\{1\},\ldots,\{n+1\},\{1,2\},\{1,2,3\},\ldots,[n+1]$$

will be a building set for P for any $n\geq 2$.

Any n-dimensional nestohedron P_B on a connected building set B can be obtained from an n-simplex by a sequence of its face truncations. In order to give the precise statement suppose $B_0 \subset B_1$ be building sets on $[n+1]$, and $S \in B_1$. Then define a *decomposition of S into elements of B_0* as $S = S_1 \sqcup \cdots \sqcup S_k$, where S_j are pairwise nonintersecting elements of B_0 and k is minimal among such disjoint representations of S. One can see easily that this decomposition exists and is unique.

Theorem 2.2 ([4, Lemma 1.5.17, Theorem 1.5.18])**.** *Every nestohedron P_B corresponding to a connected building set B can be obtained from a simplex by a sequence of face truncations.*

More precisely, let $B_0 \subset B_1$ be connected building sets on $[n+1]$. Then P_{B_1} is combinatorially equivalent to the polytope obtained from P_{B_0} by a sequence of truncations at the faces $G_i = \bigcap_{j=1}^{k_i} F_{S_j^i}$ corresponding to the decompositions $S^i = S_1^i \sqcup \cdots \sqcup S_{k_i}^i$ of elements $S^i \in B_1 \setminus B_0$, numbered in any order that is inverse to inclusion (i.e. $S^i \supset S^{i'} \Rightarrow i \leqslant i'$).

V.M.Buchstaber suggested to call a simple convex n-dimensional polytope P a *2-truncated cube* if it can be obtained from an n-cube by a sequence of cut off some faces of codimension 2 only. It is allowed to cut off any codimension 2 face that we have on a previous step of the sequence of face truncations.

Example 2.3. Here is an example of a 3-dimensional 2-truncated cube $\mathcal{P}$ which we shall use later.



FIGURE 2.1. A 2-truncated cube $\mathcal{P}$

Then any flag nestohedron can be realized as a 2-truncated cube. More precisely, the following statement holds.

Theorem 2.4 ([6])**.** *A nestohedron P_B is a flag polytope if and only if it is a 2-truncated cube.*

More precisely, if P_B is a flag polytope, then there exists a sequence of building sets $B_0 \subset B_1 \subset \cdots \subset B_N = B$, where P_{B_0} is a combinatorial cube, $B_i = B_{i-1} \cup \{S_i\}$, and P_{B_i} is obtained from $P_{B_{i-1}}$ by 2-truncation at the face $F_{S_{j_1}} \cap F_{S_{j_2}} \subset P_{B_{i-1}}$, where $S_i = S_{j_1} \sqcup S_{j_2}$, and $S_{j_1}, S_{j_2} \in B_{i-1}$.

The next family of polytopes introduced by M.Carr and S.Devadoss [7], are flag nestohedra and therefore by Theorem 2.4 can be realized as 2-truncated cubes.

Definition. *A graphical building set* $B(\Gamma)$ for a graph Γ on the vertex set $[n+1]$ consists of such S that the induced subgraph Γ_S on the vertex set $S \subset [n+1]$ is a **connected** graph.
Then $P_\Gamma = P_{B(\Gamma)}$ is called a *graph-associahedron*.

Example 2.5. The following families of graph-associahedra are of particular interest in convex geometry, combinatorics and representation theory.

- Γ is a complete graph on $[n+1]$.
  Then $P_\Gamma = Pe^n$ is a *permutohedron*, see Figure 2.2.



FIGURE 2.2. 3-dimensional permutahedron and the corresponding graph.

- Γ is a stellar graph on $[n+1]$.
  Then $P_\Gamma = St^n$ is a *stellahedron*, see Figure 2.3.



FIGURE 2.3. 3-dimensional stellahedron and the corresponding graph.

- Γ is a cycle graph on $[n+1]$.
  Then $P_\Gamma = Cy^n$ is a *cyclohedron* (or Bott-Taubes polytope [3]), see Figure 2.4.



FIGURE 2.4. 3-dimensional cyclohedron and the corresponding graph.

- Γ is a chain graph on $[n+1]$.
  Then $P_\Gamma = As^n$ is a *associahedron* (or Stasheff polytope [17]), see Figure 2.5.



FIGURE 2.5. 3-dimensional associahedron and the corresponding graph.

In order to determine the nerve-complex K_P of a graph-associahedron $P = P_\Gamma$ we should describe the combinatorial structure of its face poset. The following is a reformulation of the general property stated in [4, Theorem 1.5.13].

Proposition 2.6. *Facets of P_Γ are in 1-1 correspondence with* **non** *maximal connected subgraphs of Γ.*
Moreover, a set of facets corresponding to such subgraphs $\Gamma_{i_1}, \dots, \Gamma_{i_s}$ has a **nonempty** *intersection if and only if:*

(1) *For any two subgraphs $\Gamma_{i_k}, \Gamma_{i_l}$, either they do not have a common vertex or one is a subgraph of another;*
(2) *If any two of the subgraphs $\Gamma_{i_{k_1}}, \dots, \Gamma_{i_{k_l}}, l \geqslant 2$ do not have common vertices, then their union graph is disconnected.*

Note, that if P is a permutohedron then its facets F_1 and F_2 have a **nonempty** intersection if and only if the corresponding subgraphs Γ_1 and Γ_2 are subgraphs of one another.

3. Bigraded Betti numbers of graph-associahedra

The main result of this section generalizes [15, Theorem 2.9] to all graph-associahedra P. To compute bigraded Betti numbers $\beta^{-i,2(i+1)}(P)$ of graph-associahedra $P = P_\Gamma$ we need the following crucial statement due to A.Fenn.

Lemma 3.1 ([9, Theorem 4.6.4])**.** *Suppose P is a nestohedron, $J \subset B$. Consider the following set:*

$$\overline{J} = J \sqcup \{S \in 2^{[n+1]} \setminus B | \, \exists S_i \in J, S = S_1 \cup \ldots \cup S_k\}.$$

Then P^n_J is homotopy equivalent to $Pe^n_{\overline{J}}$.

Definition. Suppose Γ is a graph. For any of its connected subgraphs γ one can compute the number $i(\gamma)$ of such connected subraphs $\tilde{\gamma}$ in Γ that either γ and $\tilde{\gamma}$ have a nontrivial and nonempty intersection or they have no common vertices but the union of the two subgraphs is a connected subgraph in Γ.

We denote by $i_{max} = i_{max}(\Gamma)$ the maximal value of $i(\gamma)$ over all connected subgraphs γ in Γ. A connected subgraph γ, on which i_{max} is achieved, will be called a *special subgraph*.

Example 3.2. On the Figure 2.5 we have 3 special subgraphs: $\{1,2\}, \{1,4\}$ and $\{2,3\}$. The number i_{max} is equal to 4, in particular, for $\gamma = \{1,2\}$ the corresponding subgraphs $\tilde{\gamma}$ are: $\{3\}, \{4\}, \{1,4\}, \{2,3\}$.

The following statement for the bigraded Betti numbers of the type $\beta^{-i,2(i+1)}(P)$ for graph-associahedra P holds.

Theorem 3.3. *Let $P = P_\Gamma$ be a graph-associahedron of dimension $n \geq 3$ for a connected graph Γ. Then for $i > i_{max}(\Gamma)$*

$$\beta^{-i,2(i+1)}(P) = 0.$$

Denote the number of special subgraphs in Γ by s and $\omega = -i_{max}, 2(i_{max}+1)$. Then

$$\beta^{\omega}(P) = s.$$

Proof. By Theorem 1.2 and Proposition 2.6 it is sufficient to prove the following:

(a) We have: $\mathrm{cc}(P_J) \leq 2$ if $|J| > i_{max}$. Moreover, if $J = J_1 \sqcup J_2$ with $|J_{1,2}| \geq 2$, then there exists another $J' \subset [m]$ such that $J' = J'_1 \sqcup J'_2$ with $|J'_1| = 1$ and $|J'| > |J|$.

(b) Suppose $\mathrm{cc}(P_J) = 2, J = J_1 \sqcup J_2, |J| > i_{max}$. Then either $|J_1| = 1$ or $|J_2| = 1$. Moreover, if $|J| = i_{max} + 1$ then J_1 consists of a special subgraph of Γ and J_2 consists of all the i_{max} connected subgraphs in Γ that either nontrivially intersect this special subgraph or have no common vertices with it, and in the latter case their union graph is connected (this means, the corresponding facets of P do not intersect the facet corresponding to the special subgraph).

(c) Suppose $|J| > i_{max} + 1$. Then $\mathrm{cc}(P_J) = 1$.

But in the case of associahedra As^n the statement (a) follows from [15, Lemmata 2.13, 2.14], the statement (b) follows from [15, Lemmata 2.15, 2.16] and the statement (c) follows from [15, Lemma 2.17].

Applying the same technics as in [15], one can show that the statements (a), (b) and (c) (obviously, the latter one follows from (a) and (b) for any simple polytope P) hold in the permutohedron case. Moreover, using Theorem 2.2, one can obtain a permutohedron $P = Pe^n$ as a result of truncation of all faces of dimensions $0, \dots, n-2$ from an n-simplex Δ^n, and by symmetry it is evident that the special graphs γ in this case are in 1-1 correspondence with the simplices on $[\frac{n+1}{2}]$ vertices in the boundary complex $\partial\Delta^n$.

The general case can now be obtained from the permutohedron one using Lemma 3.1 and Proposition 2.6. □

As an application of Theorem 3.3 the following examples can be computed explicitly in terms of the combinatorics of the graph Γ, using induction on the polytope dimension n for combinatorial enumerations in Γ on the vertex set $[n+1]$.

Corollary 3.4. *For the 4 classical series of graph-associahedra of the dimension $n \geq 3$, one has the following values of $i_{max} = q(n)$ and s.*

$$\beta^{-q,2(q+1)}(As^n) = \begin{cases} n+3, & \text{if } n \text{ is even;} \\ \frac{n+3}{2}, & \text{if } n \text{ is odd;} \end{cases}$$
$$\beta^{-i,2(i+1)}(As^n) = 0 \quad \text{for } i \geqslant q+1,$$

where $q = q(n)$ is:

$$q = q(n) = \begin{cases} \frac{n(n+2)}{4}, & \text{if } n \text{ is even;} \\ \frac{(n+1)^2}{4}, & \text{if } n \text{ is odd.} \end{cases}$$

$$\beta^{-q,2(q+1)}(Cy^n) = \begin{cases} 2n+2, & \text{if } n \text{ is even;} \\ n+1, & \text{if } n \text{ is odd;} \end{cases}$$
$$\beta^{-i,2(i+1)}(Cy^n) = 0 \quad \text{for } i \geqslant q+1,$$

where $q = q(n)$ is:

$$q = q(n) = \begin{cases} \frac{n(n+2)-2}{2}, & \text{if } n \text{ is even;} \\ \frac{(n+1)^2-2}{2}, & \text{if } n \text{ is odd.} \end{cases}$$

$$\beta^{-q,2(q+1)}(Pe^n) = \binom{n+1}{[\frac{n+1}{2}]}$$
$$\beta^{-i,2(i+1)}(Pe^n) = 0 \quad \text{for } i \geqslant q+1,$$

where $q = q(n) = 2^{n+1} - 2^{[\frac{n+1}{2}]} - 2^{[\frac{n+2}{2}]} + 1$.

$$\beta^{-q,2(q+1)}(St^n) = \binom{n}{[\frac{n}{2}]}$$
$$\beta^{-i,2(i+1)}(St^n) = 0 \quad \text{for } i \geqslant q+1,$$

where $q = q(n) = 2^n - 2^{[\frac{n}{2}]} - 2^{[\frac{n+1}{2}]} + [\frac{n+3}{2}]$.

As graph-associahedra are flag polytopes, we can apply the previous result to studying the loop homology algebra $H_*(\Omega\mathcal{Z}_P)$ for graph-associahedra P. Namely, due to [10, Theorem 4.3] the minimal number of multiplicative generators of $H_*(\Omega\mathcal{Z}_P)$ is equal to $\sum\limits_{i=1}^{m-n}\beta^{-i,2(i+1)}(P)$. Then Theorem 3.3 gives us lower bounds for the number of multiplicative generators in the Pontryagin algebra of $\mathcal{Z}_P$.

4. Massey products

We first prove a result concerning triple Massey products of 3-dimensional classes in $H^*(\mathcal{Z}_{P_\Gamma})$, where Γ is an arbitrary (possibly disconnected) graph.

Let us state the following theorem due to G.Denham and A.Suciu which gives a combinatorial criterion for a simplicial complex K to produce a non-trivial triple Massey product of 3-dimensional classes in $H^*(\mathcal{Z}_K)$.

Theorem 4.1 ([8, Theorem 6.1.1])**.** *The following are equivalent:*

1. *There exist cohomology classes $\alpha_i \in H^3(\mathcal{Z}_K), i = 1,2,3$ for which $\langle\alpha_1,\alpha_2,\alpha_3\rangle$ is defined and non-trivial.*
2. *The underlying graph (1-skeleton) of K contains an induced subgraph isomorphic to one of the five graphs in Figure.*

Moreover, all Massey products arising in this fashion are decomposable.

Applying Theorem 4.1 to the graph-associahedra case gives us the following result.

Proposition 4.2. *There is a non-trivial triple Massey product $\langle\alpha_1,\alpha_2,\alpha_3\rangle$ of 3-dimensional cohomology classes $\alpha_i \in H^3(\mathcal{Z}_{P_\Gamma})$ for $i = 1,2,3$ if and only if there is a connected component of Γ on $m \geq 4$ vertices which is different from a complete graph K_m.*

Proof. One can check easily that the third of the five graphs in the Figure is an induced subgraph in the graph of K_P for P being any of the six 3-dimensional graph-associahedra on a connected graph with 4 vertices, except for the permutohedron. It can be seen easily that there are no 5-gons in the K_P graph in the latter case, nor is there any of the first two graphs in the Figure as an induced subgraph. The case of a connected graph on 4 vertices now holds from Theorem 4.1.

The general case follows from Proposition 2.6 and Theorem 1.1 and the dimension 3 case as if two graphs Γ_1 and Γ_2 are disjoint then for their union graph Γ one gets: $P_\Gamma = P_{\Gamma_1}\times P_{\Gamma_2}$ and the moment-angle functor $\mathcal{Z}$ preserves products of polytopes. This finishes the proof. □

Remark. Note, that each of the six 3-dimensional graph-associahedra $P = P_\Gamma$ mentioned above is a 2-truncated cube and, moreover, Pe^3 can be obtained from Cy^3 by cut off its 2 non-adjacent edges, if realized as a simplex truncation (see Theorem 2.2). As $\mathcal{Z}_P$ for $P = I^n$ is a product of spheres and therefore is a formal manifold, it follows that a nontrivial higher Massey product in $H^*(\mathcal{Z}_P)$ can either appear or vanish after a (codimension 2) face truncation (or after a stellar subdivision on the dual simplicial sphere K_P).

Example 4.3. Consider $P = Pe^3$, see Figure 2.2. It has $n = 3$ and $m = 14$, let us label its facets by the numbers $1, \dots, 14$ such that the bottom and upper 6-gon facets are 1 and 14 resp., the bottom facets are labeled by $2, \dots, 7$ and the upper facets are labeled by $8, \dots, 13$, both clockwisely.

Consider the following 3-cocycles:

$$a_1 = v_1u_{14}, a_2 = v_6u_{10}, a_3 = v_8u_4, a_4 = v_2u_{12}.$$

They correspond to 4 pairs of parallel facets of P if realized as a result of face truncations from Δ^3. Suppose they are representatives of the cohomology classes $\alpha_i \in H^3(\mathcal{Z}_P)$, that is $\alpha_i = [a_i]$ for $i = 1, \dots, 4$.

Then we get the following defining system A (see [14]) for the Massey 4-product $\langle \alpha_1, \alpha_2, \alpha_3, \alpha_4 \rangle$ (up to signs):

$$a_{13} = v_6u_1u_{14}u_{10}, a_{24} = v_6u_{10}u_8u_4, a_{35} = v_2u_8u_4u_{12},$$

$$a_{14} = v_6u_1u_8u_4u_{10}u_{14}, a_{25} = 0,$$

so $0 \in \langle \alpha_1, \alpha_2, \alpha_3, \alpha_4 \rangle$. Thus the two 3-products $\langle \alpha_1, \alpha_2, \alpha_3 \rangle$ and $\langle \alpha_2, \alpha_3, \alpha_4 \rangle$ are defined and vanish simultaneously and the 4-product $\langle \alpha_1, \alpha_2, \alpha_3, \alpha_4 \rangle$ is trivial.

We next consider a particular family of 2-truncated n-cubes $\mathcal{P}$, one for each dimension n, for which $\mathcal{Z}_\mathcal{P}$ has a nontrivial Massey product of order n.

Definition. Suppose I^n is an n-dimensional cube with facets $F_1, \dots, F_{2n}$, such that F_i and F_{n+i}, $1 \le i \le n$ are parallel (do not intersect). Then we define $\mathcal{P}$ as a result of a consecutive cut of faces of codimension 2 from I^n, having the following Stanley-Reisner ideal:

$$I = (v_1v_{n+1}, \dots, v_nv_{2n}, v_1v_{n+2}, \dots, v_{n-1}v_{2n}, \dots, v_1v_{2n-1}, v_2v_{2n}, \dots),$$

where v_i correspond to F_i, $1 \le i \le 2n$ and in the last dots are the monomials corresponding to the new facets. This determines uniquely the combinatorial type of $\mathcal{P}$.

Remark. (1) Note that for $n = 2$ we get a 2-dimensional cube (the square) and for $n = 3$ we get a simple polytope P for which $K = K_P$ is a simplicial complex with a nontrivial triple Massey product in $H^*(\mathcal{Z}_K)$ due to the result of Baskakov, see [2].

(2) The 2-truncated cube $\mathcal{P}$ is not a graph-associahedron, due to [6, Theorem 9.2], as its number of facets $f_0(\mathcal{P}) = \frac{n(n+3)}{2} - 1 < f_0(As^n) = \frac{n(n+3)}{2}$. However, we can easily construct the building set B for $\mathcal{P}$ on the vertex set $S = [n+1]$ by identifying F_i with $\{1, \dots, i\}$ for $1 \le i \le n$ and identifying F_i with $\{i-n+1\}$ for $n+1 \le i \le 2n$. Then by Theorem 2.4, we consecutively cut the following faces:

$$\{1\} \sqcup \{3\}, \{1,2\} \sqcup \{4\}, \dots, \{1, \dots, n-1\} \sqcup \{n+1\}$$

$$\dots$$

$$\{1\} \sqcup \{n\}, \{1,2\} \sqcup \{n+1\}.$$

Thus, $\mathcal{P} = P_B$ for the the building set B consisting of the building set B_0 of an n-cube from Example 2.1, the above subsets in $[n+1]$ and all the subsets in $[n+1]$ which are the unions of nontrivially intersecting elements in B.

Theorem 4.4. *Let $\alpha_i \in H^3(\mathcal{Z}_\mathcal{P})$ be represented by a 3-cocycle v_iu_{n+i} for $1 \leq i \leq n$ and $n \geq 2$. Then all Massey products of consecutive elements from $\alpha_1, \ldots, \alpha_n$ are defined and the whole n-product $\langle \alpha_1, \ldots, \alpha_n \rangle$ is nontrivial.*

Proof. Let us prove the theorem by induction on n. The base case $n = 2$ is trivial: α_1 and α_2 are the classes of 3-dimensional spheres in $\mathcal{Z}_\mathcal{P} \cong S^3 \times S^3$ and their cup-product (i.e Massey 2-product) is the dual to the fundamental class of $\mathcal{Z}_\mathcal{P}$.

We first note that all Massey products of orders less than n vanish simultaneously in $H^{*,*}(\mathcal{Z}_\mathcal{P}) \cong H\big[\Lambda[u_1, \ldots, u_m] \otimes \Bbbk[\mathcal{P}], d\big]$, i.e contain coboundaries. Starting with the representing cocycles v_iu_{i+n} of α_i it can be seen by induction on the dimension n of $\mathcal{P}$ that if a defining system C for the matrix n-product $\langle \alpha_1, \ldots, \alpha_n \rangle$ can be extended from ith diagonal of the matrix C to its $(i+1)$th diagonal for all $2 \leq i \leq n$ then $c_{lm}, m - l = i \geq 2$ have either a form $v_ku_{j_1} \ldots u_{j_{2i-1}}$ or a form $v_ku_{j_1} \ldots u_{j_{2i-1}} + d(u_ku_{j_1} \ldots u_{j_{2i-1}})$ (up to the signs). The latter can be checked as the differential in the cohomology algebra preserves multigrading and by using the codimension 2 face cuts from the definition of $\mathcal{P}$ (see also the Example below).

Then the Massey n-product $\langle \alpha_1, \ldots, \alpha_n \rangle$ is defined and any cohomology class, belonging to it, lies in the multigraded component $H^{-(2n-2),(2,\ldots,2)}(\mathcal{Z}_\mathcal{P})$ of the moment-angle manifold $\mathcal{Z}_\mathcal{P}$ with one of its representatives being the class of the cocycle $v_1v_{2n}u_2 \ldots u_{2n-1}$. Up to sign we have the following equality for any representative c of an element in $\langle \alpha_1, \ldots, \alpha_n \rangle$ for any defining system C, see [14]:

$$c = d(c_{1,n+1}) - (-1)^3 v_1u_{n+1}c_{2,n+1} - \overline{c}_{1,3}c_{3,n+1} - \ldots - \overline{c}_{1,n}v_nu_{2n},$$

where $(n+1)\times(n+1)$-matrix C is upper triangular with zeros on the diagonal and $c_{i,i+1} = -v_iu_{n+i}$ for $1 \leq i \leq n$, such that the following condition holds:

$$cE_{1,n+1} = d(C) - \overline{C} \cdot C$$

and $\overline{c}_{ij} = (-1)^{|c_{ij}|}c_{ij}$ depends on the degree $|c_{ij}|$ of a matrix element c_{ij}.

By definition of higher Massey operations (see [14]), one has: $d(c_{2,n+1})$ is a representative in $\langle \alpha_2, \ldots, \alpha_n \rangle$ and $d(c_{1,n})$ is a representative in $\langle \alpha_1, \ldots, \alpha_{n-1} \rangle$. To prove that $v_1v_{2n}u_2 \ldots u_{2n-1}$ is the only representing cocycle for the n-product we use induction on n, the represnting monomials for the indeterminacies and the multigrading in $H^*(\mathcal{Z}_\mathcal{P})$, see Theorem 1.3. For instance, the indeterminacy for the first of the $(n-1)$-products above lies in the multigraded component of $v_2u_3 \ldots u_nu_{n+2} \ldots u_{2n}$ and the only cocycle there is the coboundary $d(u_2 \ldots u_nu_{n+2} \ldots u_{2n})$. As for the second $(n-1)$-product above, the indeterminacy lies in the multigraded component of $v_1u_2 \ldots u_{n-1}u_{n+1} \ldots u_{2n-1}$ and the only cocycle there is the coboundary $d(u_1 \ldots u_{n-1}u_{n+1} \ldots u_{2n-1})$.

Thus, the Massey n-product $\langle \alpha_1, \ldots, \alpha_n \rangle$ is defined and nontrivial, consisting only of the cohomology class of $v_1v_{2n}u_2 \ldots u_{2n-1}$. □

Remark. Note, that the nontrivial n-product constructed above is decomposable. Namely, one has: $[v_1v_{2n}u_2 \ldots u_{2n-1}] = \pm[v_1u_{n+1} \ldots u_{2n-1}] \cdot [v_{2n}u_2 \ldots u_n]$.

Example 4.5. Consider the case $n = 4$. Then the Stanley–Reisner ideal of $\mathcal{P}$ is

$$I = (v_1v_5, v_2v_6, v_3v_7, v_4v_8, v_1v_6, v_2v_7, v_3v_8, v_1v_7, v_2v_8, \dots)$$

and the cohomology classes $\alpha_i, 1 \le i \le 4$ are represented by the cocycles $a_i = v_iu_{4+i}, 1 \le i \le 4$. One has (up to sign):

$$a_1a_2 = d(v_1u_2u_5u_6), a_2a_3 = d(v_2u_3u_6u_7), a_3a_4 = d(v_2u_4u_7u_8).$$

Alternatively, one has (up to sign):

$$a_1a_2 = d(v_2u_1u_5u_6 - v_5u_2u_1u_6 + v_6u_2u_1u_5),$$

$$a_2a_3 = d(v_3u_2u_6u_7 - v_6u_3u_2u_7 + v_7u_3u_2u_6),$$

$$a_3a_4 = d(v_4u_2u_7u_8 - v_7u_4u_2u_8 + v_8u_4u_2u_7).$$

Then in first case one has the following cocycle representing a class in $\langle\alpha_1, \alpha_2, \alpha_3\rangle$ (here the Massey 2-product of a and b is equal to $\overline{a} \cdot b$, $\overline{a} = (-1)^{|a|}a$):

$$v_1u_5 \cdot (-v_2u_3u_6u_7) - v_1u_2u_5u_6 \cdot v_3u_7 = d(v_1u_2u_3u_5u_6u_7)$$

and the following cocycle representing a class in $\langle\alpha_2, \alpha_3, \alpha_4\rangle$:

$$v_2u_6 \cdot (-v_3u_4u_7u_8) - v_2u_3u_6u_7 \cdot v_4u_8 = d(v_2u_3u_4u_6u_7u_8).$$

Similarly, in the second case the representing cocycle for $\langle\alpha_1, \alpha_2, \alpha_3\rangle$ will be $d(v_3u_1u_2u_5u_6u_7)$ and for $\langle\alpha_2, \alpha_3, \alpha_4\rangle$ one gets: $d(v_4u_2u_3u_6u_7u_8)$.

Thus, the Massey products $\langle\alpha_1, \alpha_2, \alpha_3\rangle$ and $\langle\alpha_2, \alpha_3, \alpha_4\rangle$ vanish simultaneously in both cases and the 4-product $\langle\alpha_1, \alpha_2, \alpha_3, \alpha_4\rangle$ is defined. Considering the multigrading in $H^*(\mathcal{Z}_P)$ it is easy to see that the latter 4-product consists of the only class with a representative $v_1v_8u_2 \dots u_7$ in $H^{-6,(2,\dots,2)}(\mathcal{Z}_P) \subset H^{-6,16}(\mathcal{Z}_P) \subset H^{10}(\mathcal{Z}_P)$, where $\mathcal{Z}_P$ is a closed smooth 17-dimensional manifold. More precisely, the representing cocycle c for $\langle\alpha_1, \alpha_2, \alpha_3, \alpha_4\rangle$ is equal to:

$$d(c_{1,5}) - \overline{a}_1c_{2,5} - \overline{c}_{1,3}c_{3,5} - \overline{c}_{1,4}a_4$$

which is equal to

$$d(c_{1,5}) - d(v_1u_2 \dots u_8) + v_1v_8u_2 \dots u_7$$

in the first case and to (up to signs)

$$d(c_{1,5}) + d(v_4 \dots) + d(v_7 \dots) + d(v_8 \dots) + v_1v_8u_2 \dots u_7$$

in the second case. Other two possible cases for $c_{2,5}$ and $c_{1,4}$ (see Theorem 4.4) give the same representing cocycle for the class c.

Finally, one has: $[v_1v_8u_2 \dots u_7] = -[v_1u_5u_6u_7] \cdot [v_8u_2u_3u_4]$.

Using Theorem 4.4 we can construct a smooth closed 2-connected manifold M with a compact torus action, such that there are nontrivial higher Massey products of any prescribed orders $n_1, \dots, n_r$ in $H^*(M)$. Namely, we let $M = \mathcal{Z}_P$, where $P = \mathcal{P}^{n_1} \times \dots \times \mathcal{P}^{n_r}$ and remind that the functor $\mathcal{Z}$ preserves products for simple polytopes. Note that P is still a flag polytope but is no more a 2-truncated cube, in general.

References

[1] Anthony Bahri, Martin Bendersky, Frederick R. Cohen, and Samuel Gitler. *The polyhedral product functor: a method of computation for moment-angle complexes, arrangements and related spaces*, Adv. Math. **225** (2010), no. 3, 1634–1668.

[2] I. V. Baskakov. *Massey triple products in the cohomology of moment-angle complexes*, Russian Math. Surveys, 58:5 (2003), 1039–1041.

[3] Raul Bott and Clifford Taubes. *On the self-linking of knots. Topology and physics.* J.Math.Phys. **35** (1994), no. 10, 5247–5287.

[4] Victor M. Buchstaber and Taras E. Panov. *Toric Topology*, Mathematical Surveys and Monographs, 204, American Mathematical Society, Providence, RI, 2015.

[5] Victor M. Buchstaber and Taras E. Panov. *Torus actions, combinatorial topology and homological algebra.* Uspekhi Mat. Nauk **55** (2000), no. 5, 3–106 (Russian). Russian Math. Surveys **55** (2000), no. 5, 825–921 (English translation).

[6] Victor M. Buchstaber, Vadim Volodin. *Precise upper and lower bounds for nestohedra.* Izv. Ross. Akad. Nauk, Ser. Mat. **75** (2011), no. 6, 17–46 (Russian); Izv. Math. **75** (2011), no. 6 (English translation).

[7] Michael P.Carr and Satyan L.Devadoss. *Coxeter complexes and graph-associahedra*, Topology Appl. **153** (2006), no. 12, 2155–2168.

[8] Graham Denham and Alexander I. Suciu. *Moment-angle complexes, monomial ideals, and Massey products*, Pure and Applied Mathematics Quarterly, 3(1) (2007), (Robert MacPherson special issue, part 3), 25–60.

[9] A. Fenn, *On families of nestohedra*, PhD thesis, Manchester University, 2010.

[10] Jelena Grbic, Taras Panov, Stephen Theriault and Jie Wu. *Homotopy types of moment-angle complexes for flag complexes*, Transactions of the AMS, (2015), to appear; arXiv:1211.0873.

[11] Jelena Grbić and Stephen Theriault. *The homotopy type of the complement of a coordinate subspace arrangement*, Topology **46** (2007), no. 4, 357–396.

[12] M. Hochster, *Cohen-Macaulay rings, combinatorics, and simplicial complexes*, in Ring theory, II (Proc. Second Conf.,Univ. Oklahoma, Norman, Okla., 1975), Lecture Notes in Pure and Appl. Math., V. 26, 171–223, Dekker, New York, 1977.

[13] Kouyemon Iriye and Daisuke Kishimoto. *Decompositions of polyhedral products for shifted complexes*, Adv. Math. **245** (2013), 716–736.

[14] David Kraines, *Massey higher products*, Trans. Amer. Math. Soc., **124** (1966), 431–449.

[15] Ivan Yu. Limonchenko, *Bigraded Betti numbers of certain simple polytopes*, Mathematical Notes, 94:3 (2013), 351-363.

[16] Taras E. Panov. *Cohomology of face rings, and torus actions*, in "Surveys in Contemporary Mathematics". London Math. Soc. Lecture Note Series, vol. **347**, Cambridge, U.K., 2008, 165–201; arXiv:math.AT/0506526.

[17] James D. Stasheff. *Homotopy associativity of H-spaces. I.* Transactions Amer. Math. Soc. **108** (1963), 275–292.

Department of Geometry and Topology, Faculty of Mathematics and Mechanics, Moscow State University, Leninskiye Gory, Moscow 119992, Russia

E-mail address: ilimonchenko@gmail.com